

**APPENDIX**

Versions of amended title, specification, and claims 1, 28, 29, 30, 33, 55, and 56, with markings to show changes made, pursuant to 37 C.F.R. § 1.121 (c)(1)(ii):

**AMENDED TITLE**

Applicant presents the amended title in marked-up form below, to aid the Examiner in identifying amendments.

DYE COMPOSITION CONTAINING 1,8-BIS(2,5-DIAMINOPHENOXY)-3,[5]6-DIOXAOCANE, AN ADDITIONAL OXIDATION BASE AND A COUPLER, AND DYEING PROCESSES

**AMENDED SPECIFICATION**

Applicant presents the amended portions of the specification in marked-up form below, to aid the Examiner in identifying amendments.

*Amended first full paragraph on page 1 of the specification:*

The invention relates to a composition for the oxidation dyeing of keratin fibers, containing a first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaocane, and the acid-addition salts thereof, at least one second selected oxidation base and at least one coupler; as well as to the oxidation dyeing process using this composition.

*Amended last full paragraph on page 2, which extends to the first two lines of page 3, of the specification:*

The inventor has now discovered, entirely surprisingly and unexpectedly, that the combination of 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaocane, and/or of at least one of the acid-addition salts thereof, with at least one second suitably selected oxidation base and at least one coupler, can give intense colorations which moreover can have improved properties of

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

resistance with respect to the various attacking factors to which the hair may be subjected (shampooing, light, bad weather, permanent-waving, perspiration, friction, etc.).

*Amended paragraph on page 3, which extends from line 6 to line 7, of the specification:*

- at least one first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane and acid-addition salts thereof,

*Amended paragraph on page 8, which extends from line 6 to line 9, of the specification:*

The at least one first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane and acid-addition salts thereof preferably represent from 0.0005 to 12% by weight approximately relative to the total weight of the dye composition, and even more preferably from 0.005 to 6% by weight approximately relative to this weight.

*Amended table on page 15 of the specification:*

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

EXAMPLE	1	2	3	4	5	6
1,8-Bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane tetrahydrochloride monohydrate	0.39	0.39	0.39	0.39	0.39	0.39
para-Phenylenediamine (second oxidation base)	0.162	-	0.162	-	-	-
para-Aminophenol (second oxidation base)	-	0.163	-	-	-	-
3-Methyl-4-aminophenol (second oxidation base)	-	-	-	0.184	-	-
2-( $\beta$ -Hydroxyethyl-para-phenylenediamine dihydrochloride (second oxidation base)	-	-	-	-	0.337	-
2,6-Dimethyl-para-phenylenediamine dihydrochloride (second oxidation base)	-	-	-	-	-	0.313
5-N-( $\beta$ -Hydroxyethyl)amino-2-methylphenol (coupler)	0.498	-	-	-	-	-
2,4-Diaminophenoxyethanol dihydrochloride (coupler)	-	0.723	-	-	-	-
1,3-Dihydroxybenzene (coupler)	-	-	0.33	-	-	-
5-Amino-2-methylphenol (coupler)	-	-	-	-	-	-
3-Aminophenol	-	-	-	-	0.327	-
6-Hydroxybenzomorpholine	-	-	-	-	-	0.453

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

*Amended table on page 19 of the specification:*


EXAMPLE	7	8	9	10	11	12
1,8-Bis(2,5-diaminophenoxy)-3,5-bis(2,5-dioxaoctane tetra hydrochloride monohydrate	0.39	0.39	0.39	0.39	0.39	0.39
para-Phenylenediamine (second oxidation base)	0.162	-	0.162	-	-	-
para-Aminophenol (second oxidation base)	-	0.163	-	-	-	-
3-Methyl-4-aminophenol (second oxidation base)	-	-	-	0.184	-	-
2-(β-Hydroxyethyl)-para-phenylenediamine dihydrochloride (second oxidation base)	-	-	-	-	0.337	-
2,6-Dimethyl-para-phenylenediamine dihydrochloride (second oxidation base)	-	-	-	-	-	0.313
5-N-(β-Hydroxyethyl)amino-2-methylphenol (coupler)	0.498	-	-	-	-	-
2,4-Diaminophenoxyethanol dihydrochloride (coupler)	-	0.723	-	-	-	-
1,3-Dihydroxybenzene (coupler)	-	-	0.33	-	-	-
5-Amino-2-methylphenol (coupler)	-	-	-	0.369	-	-
3-Aminophenol	-	-	-	-	0.327	-
6-Hydroxybenzomorpholine	-	-	-	-	-	0.453

**AMENDED CLAIMS**

Applicant presents amended claims 1, 28-30, 33, 55 and 56 in marked-up form below, to aid the Examiner in identifying amendments.

1. (Amended) A composition for oxidation dyeing of keratin fibers, comprising:

- at least one first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,5-dioxaoctane and acid-addition salts thereof;
- at least one second oxidation base chosen from para-phenylenediamine, para-toluenediamine, N,N-bis-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, 2-chloro-para-phenylenediamine, N-phenyl-para-phenylenediamine, 4,4'-diaminodiphenylamine, N-methoxyethyl-para-phenylenediamine, 2-n-propyl-para-phenylenediamine, 4-aminophenol, N-methyl-4-aminophenol, 2-hydroxymethyl-4-aminophenol, 3-methyl-4-aminophenol, 2-aminomethyl-4-aminophenol, 2-( $\beta$ -hydroxyethylaminomethyl)-4-aminophenol, 2-methoxy-4-aminophenol, 2-methoxymethyl-4-aminophenol, tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-N-methylpyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, N,N'-bis( $\beta$ -hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, 3-amino-6-dimethylaminopyridine and pyrazolo[1,5-a]pyrimidines of formula (I):



in which:

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

- R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>, which are identical or different, are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals wherein said amino can be protected with a protective group chosen from acetyl, ureido and sulphonyl groups, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, and di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino-(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals;

- radicals X are identical or different, and are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, amino radicals, (C<sub>1</sub>-C<sub>4</sub>)alkyl-amino radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino radicals, halogen atoms, carboxylic acid groups and sulphonic acid groups;

- i is chosen from 0, 1, 2 and 3;

- p is chosen from 0 and 1;

- q is chosen from 0 and 1;

- n is chosen from 0 and 1;

with the proviso that:

- (i) the sum  $p + q$  is other than 0;

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

- (ii) when  $p + q$  is equal to 2, then  $n$  is 0 and the groups  $NR_1R_2$  and  $NR_3R_4$  occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
  - (iii) when  $p$  is equal to 1 and  $q$  is equal to 0, then  $n$  is 1 and the group  $NR_1R_2$  and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
  - (iv) when  $p$  is equal to 0 and  $q$  is equal to 1, then  $n$  is 1 and the group  $NR_3R_4$  and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
- and acid-addition salts thereof;
- and at least one coupler.

28. (Amended) A composition for oxidation dyeing of keratin fibers comprising: 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane tetrahydrochloride monohydrate, para-phenylenediamine, 5- N-( $\beta$ -hydroxyethyl)amino-2-methylphenol, ethanol, sodium metabisulphite, pentasodium diethylenetriaminepentaacetic acid, aqueous ammonia, and demineralized water.

29. (Amended) A composition for oxidation dyeing of keratin fibers comprising: 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane tetrahydrochloride monohydrate, para-phenylenediamine, 5-N-( $\beta$ -hydroxyethyl)amino-2-methylphenol, ethanol, dipotassium hydrogenphosphate, potassium dihydrogenphosphate, sodium metabisulphite, pentasodium diethylenetriaminepentaacetic acid, and demineralized water.

30. (Amended) A composition for oxidation dyeing of keratin fibers comprising


- at least one oxidation base chosen from acid-addition salts of 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane, wherein said salts are chosen from hydrochlorides, hydrobromides, sulphates, citrates, succinates, tartrates, lactates and acetates;

- at least one second oxidation base chosen from

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

para-phenylenediamine, para-toluenediamine, N,N-bis-(β-hydroxyethyl)-para-phenylenediamine, 2-(β-hydroxyethyl)-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, 2-chloro-para-phenylenediamine, N-phenyl-para-phenylenediamine, 4,4-diaminodiphenylamine, N-methoxyethyl-para-phenylenediamine, 2-n-propyl-para-phenylenediamine, 4-aminophenol, N-methyl-4-aminophenol, 2-hydroxymethyl-4-aminophenol, 3-methyl-4-aminophenol, 2-aminomethyl-4-aminophenol, 2-(β-hydroxyethylaminomethyl)-4-aminophenol, 2-methoxy-4-aminophenol, 2-methoxymethyl-4-aminophenol, tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-N-methylpyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, N,N'-bis(β-hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, 3-amino-6-dimethylaminopyridine and pyrazolo[1,5-a]pyrimidines of formula (I):



in which:

- R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>, which are identical or different, are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals wherein said amino can be protected with a protective group chosen from acetyl, ureido and sulphonyl groups, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings,

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com



hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, and di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino-(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals;

- radicals X are identical or different, and are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals,

di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings,

hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, amino radicals, (C<sub>1</sub>-C<sub>4</sub>)alkyl-amino radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino radicals, halogen atoms, carboxylic acid groups and sulphonic acid groups;

- i is chosen from 0, 1, 2 and 3;

- p is chosen from 0 and 1;

- q is chosen from 0 and 1;

- n is chosen from 0 and 1;

with the proviso that:

- (i) the sum  $p + q$  is other than 0;

- (ii) when  $p + q$  is equal to 2, then n is 0 and the groups NR<sub>1</sub>R<sub>2</sub> and NR<sub>3</sub>R<sub>4</sub> occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

- (iii) when p is equal to 1 and q is equal to 0, then n is 1 and the group NR<sub>1</sub>R<sub>2</sub> and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

- (iv) when p is equal to 0 and q is equal to 1, then n is 1 and the group NR<sub>3</sub>R<sub>4</sub> and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

and acid-addition salts thereof;

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

- and at least one coupler chosen from meta-phenylenediamines, meta-aminophenols, meta-diphenols, heterocyclic couplers, sesamol,  $\alpha$ -naphthol, and acid-addition salts thereof.

33. (Amended) A process for oxidation dyeing of keratin fibers, comprising:

applying to keratin fibers to be dyed a dyeing composition;

developing a desired color in said keratin fibers with the aid of at least one oxidizing agent;


wherein said dyeing composition comprises:

- at least one first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,5-dioxaoctane and acid-addition salts thereof,
- at least one second oxidation base chosen from para-phenylenediamine, para-toluenediamine, N,N-bis-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, 2-chloro-para-phenylenediamine, N-phenyl-para-phenylenediamine, 4,4'-diaminodiphenylamine, N-methoxyethyl-para-phenylenediamine, 2-n-propyl-para-phenylenediamine, 4-aminophenol, N-methyl-4-aminophenol, 2-hydroxymethyl-4-aminophenol, 3-methyl-4-aminophenol, 2-aminomethyl-4-aminophenol, 2-( $\beta$ -hydroxyethylaminomethyl)-4-aminophenol, 2-methoxy-4-aminophenol, 2-methoxymethyl-4-aminophenol, tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-N-methylpyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, N,N'-bis( $\beta$ -hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, 3-amino-6-dimethylaminopyridine and pyrazolo[1,5-a]pyrimidines of formula

(I):

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com



in which:

- R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>, which are identical or different, are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals wherein said amino can be protected with a protective group chosen from acetyl, ureido and sulphonyl groups, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, and di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino-(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals;
- radicals X are identical or different, and are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, amino radicals, (C<sub>1</sub>-C<sub>4</sub>)alkyl-amino radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino radicals, halogen atoms, carboxylic acid groups and sulphonic acid groups;
- i is chosen from 0, 1, 2 and 3;

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

- p is chosen from 0 and 1;

- q is chosen from 0 and 1;

- n is chosen from 0 and 1;

with the proviso that:

- (i) the sum  $p + q$  is other than 0;
- (ii) when  $p + q$  is equal to 2, then n is 0 and the groups  $NR_1R_2$  and  $NR_3R_4$  occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
- (iii) when p is equal to 1 and q is equal to 0, then n is 1 and the group  $NR_1R_2$  and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
- (iv) when p is equal to 0 and q is equal to 1, then n is 1 and the group  $NR_3R_4$  and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

and acid-addition salts thereof;

- and at least one coupler.

55. (Amended) A multi-compartment dyeing device, comprising:

a first compartment,

a second compartment;


wherein said first compartment contains a dyeing composition comprising:

- at least one first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,[5]6-dioxaoctane and acid-addition salts thereof;
- at least one second oxidation base chosen from para-phenylenediamine, para-toluenediamine, N,N-bis-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, 2-chloro-para-phenylenediamine, N-phenyl-para-phenylenediamine, 4,4-diaminodiphenylamine, N-

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

methoxyethyl-para-phenylenediamine, 2-n-propyl-para-phenylenediamine, 4-aminophenol, N-methyl-4-aminophenol, 2-hydroxymethyl-4-aminophenol, 3-methyl-4-aminophenol, 2-aminomethyl-4-aminophenol, 2-(β-hydroxyethylaminomethyl)-4-aminophenol, 2-methoxy-4-aminophenol, 2-methoxymethyl-4-aminophenol, tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-N-methylpyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, N,N'-bis(β-hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, 3-amino-6-dimethylaminopyridine and pyrazolo[1,5-a]pyrimidines of formula (I):



in which:

- R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub> and R<sub>4</sub>, which are identical or different, are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkoxy(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals wherein said amino can be protected with a protective group chosen from acetyl, ureido and sulphonyl groups, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, and di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino-(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals;

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

- radicals X are identical or different, and are chosen from a hydrogen atom, C<sub>1</sub>-C<sub>4</sub> alkyl radicals, aryl radicals, C<sub>1</sub>-C<sub>4</sub> hydroxyalkyl radicals, C<sub>2</sub>-C<sub>4</sub> polyhydroxyalkyl radicals, C<sub>1</sub>-C<sub>4</sub> aminoalkyl radicals, (C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkylamino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, di[hydroxy(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino(C<sub>1</sub>-C<sub>4</sub>)alkyl radicals, amino radicals, (C<sub>1</sub>-C<sub>4</sub>)alkyl-amino radicals, di[(C<sub>1</sub>-C<sub>4</sub>)alkyl]amino radicals, halogen atoms, carboxylic acid groups and sulphonic acid groups;

- i is chosen from 0, 1, 2 and 3;

- p is chosen from 0 and 1;

- q is chosen from 0 and 1;

- n is chosen from 0 and 1;

with the proviso that:

- (i) the sum  $p + q$  is other than 0;

- (ii) when  $p + q$  is equal to 2, then n is 0 and the groups NR<sub>1</sub>R<sub>2</sub> and NR<sub>3</sub>R<sub>4</sub> occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

- (iii) when p is equal to 1 and q is equal to 0, then n is 1 and the group NR<sub>1</sub>R<sub>2</sub> and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

- (iv) when p is equal to 0 and q is equal to 1, then n is 1 and the group NR<sub>3</sub>R<sub>4</sub> and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

and acid-addition salts thereof;

- and at least one coupler;

wherein said second compartment contains an oxidizing composition comprising:

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

- at least one oxidizing agent.


56. (Amended) A dyeing kit comprising:

a first container,

a second container;

wherein said first container contains a dyeing composition comprising:

- at least one first oxidation base chosen from 1,8-bis(2,5-diaminophenoxy)-3,5-dioxaoctane and acid-addition salts thereof;
- at least one second oxidation base chosen from para-phenylenediamine, para-toluenediamine, N,N-bis-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2-( $\beta$ -hydroxyethyl)-para-phenylenediamine, 2,6-dimethyl-para-phenylenediamine, 2-isopropyl-para-phenylenediamine, 2-chloro-para-phenylenediamine, N-phenyl-para-phenylenediamine, 4,4-diaminodiphenylamine, N-methoxyethyl-para-phenylenediamine, 2-n-propyl-para-phenylenediamine, 4-aminophenol, N-methyl-4-aminophenol, 2-hydroxymethyl-4-aminophenol, 3-methyl-4-aminophenol, 2-aminomethyl-4-aminophenol, 2-( $\beta$ -hydroxyethylaminomethyl)-4-aminophenol, 2-methoxy-4-aminophenol, 2-methoxymethyl-4-aminophenol, tetraaminopyrimidine, 4-hydroxy-2,5,6-triaminopyrimidine, 4,5-diamino-1-ethyl-3-methylpyrazole, 4,5-diamino-N-methylpyrazole, 4,5-diamino-1-(4'-chlorobenzyl)pyrazole, N,N'-bis( $\beta$ -hydroxyethyl)-N,N'-bis(4'-aminophenyl)-1,3-diaminopropanol, 3-amino-6-dimethylaminopyridine and pyrazolo[1,5-a]pyrimidines of formula (I):



FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

in which:

- $R_1$ ,  $R_2$ ,  $R_3$  and  $R_4$ , which are identical or different, are chosen from a hydrogen atom,  $C_1$ - $C_4$  alkyl radicals, aryl radicals,  $C_1$ - $C_4$  hydroxyalkyl radicals,  $C_2$ - $C_4$  polyhydroxyalkyl radicals,  $(C_1$ - $C_4$ )alkoxy( $C_1$ - $C_4$ )alkyl radicals,  $C_1$ - $C_4$  aminoalkyl radicals wherein said amino can be protected with a protective group chosen from acetyl, ureido and sulphonyl groups,  $(C_1$ - $C_4$ )alkylamino( $C_1$ - $C_4$ )alkyl radicals, di[( $C_1$ - $C_4$ )alkyl]amino( $C_1$ - $C_4$ )alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy( $C_1$ - $C_4$ )alkylamino( $C_1$ - $C_4$ )alkyl radicals, and di[hydroxy( $C_1$ - $C_4$ )alkyl]amino-( $C_1$ - $C_4$ )alkyl radicals;
- radicals X are identical or different, and are chosen from a hydrogen atom,  $C_1$ - $C_4$  alkyl radicals, aryl radicals,  $C_1$ - $C_4$  hydroxyalkyl radicals,  $C_2$ - $C_4$  polyhydroxyalkyl radicals,  $C_1$ - $C_4$  aminoalkyl radicals,  $(C_1$ - $C_4$ )alkylamino( $C_1$ - $C_4$ )alkyl radicals, di[( $C_1$ - $C_4$ )alkyl]amino( $C_1$ - $C_4$ )alkyl radicals, wherein said dialkyls can form a ring chosen from 5- and 6-membered aliphatic and heterocyclic rings, hydroxy( $C_1$ - $C_4$ )alkylamino( $C_1$ - $C_4$ )alkyl radicals, di[hydroxy( $C_1$ - $C_4$ )alkyl]amino( $C_1$ - $C_4$ )alkyl radicals, amino radicals,  $(C_1$ - $C_4$ )alkyl-amino radicals, di[( $C_1$ - $C_4$ )alkyl]amino radicals, halogen atoms, carboxylic acid groups and sulphonic acid groups;
- i is chosen from 0, 1, 2 and 3;
- p is chosen from 0 and 1;
- q is chosen from 0 and 1;
- n is chosen from 0 and 1;

with the proviso that:

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com



- (i) the sum  $p + q$  is other than 0;
- (ii) when  $p + q$  is equal to 2, then  $n$  is 0 and the groups  $NR_1R_2$  and  $NR_3R_4$  occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
- (iii) when  $p$  is equal to 1 and  $q$  is equal to 0, then  $n$  is 1 and the group  $NR_1R_2$  and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);
- (iv) when  $p$  is equal to 0 and  $q$  is equal to 1, then  $n$  is 1 and the group  $NR_3R_4$  and the OH group occupy positions (2,3); (5,6); (6,7); (3,5) and (3,7);

and acid-addition salts thereof;

- and at least one coupler;

wherein said second container contains an oxidizing composition comprising:

- at least one oxidizing agent.

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com